



3585 BROADWAY • (541) 756-2584
1-800-422-9393
FAX (541) 756-5783
NORTH BEND, OREGON • 97459

JON C. KINTNER, M.D.
DENNIS E. OTTEMILLER, M.D.
DEBRA A. GRAHAM, M.D.
JANE GILBERT, M.D.
EYE M.D.s

Practice Limited to the Eye
Cataract Surgery & Intraocular Lenses
Board Certified, Members of American
Academy of Ophthalmology

Cataracts in Children

A cataract is a clouding of the eye's normally clear lens. The lens of the eye plays an important role in focusing images on the retina, the light-sensitive nerve cells lining the back of the eye. If the lens loses its clarity, light rays do not focus clearly and vision is blurry. Just as it is hard to see through a dirty window, it is hard to see through a cataract. Although most cataracts occur in older adults, they can appear in children, in one or both eyes, often at birth. They look like a white or gray spot in the pupil.

Cataracts in children may be inherited or develop because of an infection or a disease acquired before birth, or as a result of an injury. In most cases, no specific cause is found.

Children may lose vision permanently because of amblyopia (lazy eye) if a significant cataract is not removed quickly. Mild cataracts may not need treatment.

The focusing power of the original lens, removed during cataract surgery, must be replaced to restore vision. Intraocular lenses (IOLs), permanent plastic lenses placed inside the eye, are implanted in older children much as they are in adults. In infants, IOLs are controversial because the eyes grow and change their prescriptions during the first few years of life. Many surgeons prefer contact lenses or even glasses for younger children.

Regardless of the type of correction, children need follow-up exams to avoid possible complications, including glaucoma, scar tissue forming in the pupil, and amblyopia. Often, children will need eye muscle surgery because the eye turns or crosses.