



3585 BROADWAY • (541) 756-2584  
1-800-422-9393  
FAX (541) 756-5783  
NORTH BEND, OREGON • 97459

JON C. KINTNER, M.D.  
DENNIS E. OTTEMILLER, M.D.  
DEBRA A. GRAHAM, M.D.  
JANE GILBERT, M.D.  
EYE M.D.s

Practice Limited to the Eye  
Cataract Surgery & Intraocular Lenses  
Board Certified, Members of American  
Academy of Ophthalmology

## Chalazion

A chalazion is a swelling in the eyelid caused by inflammation of one of the small oil producing glands located in the upper and lower eyelids. A chalazion is sometimes confused with a sty, which also appears as a lump in the eyelid, but is an infection of a lash follicle that forms a red, sore lump. Chalazions tend to occur farther from the edge of the eyelid than styes and tend to “point” toward the inside of the eyelid. Sometimes a chalazion can cause the entire eyelid to swell suddenly, but usually there is a definite tender point.

When a chalazion is small and without symptoms, it may disappear on its own. If the chalazion is large, it may cause blurred vision. Chalazions are treated with any or a combination of the following methods:

- Warm compresses can be applied. The simplest way is to hold a clean washcloth, soaked in hot water, against the closed lid. Do this for five to ten minutes, three or four times a day. Repeatedly soak the washcloth in hot water to maintain adequate heat. The majority of chalazions will disappear within a few weeks. Sometimes antibiotic ointments are used in combination with warm compresses.
- Surgical incision or excision may be used to remove large chalazions which do not respond to other treatments.

Chalazions usually respond well to treatment, although some people are prone to recurrences. If a chalazion recurs in the same place, your ophthalmologist (Eye M.D.) may suggest a biopsy to rule out more