



3585 BROADWAY • (541) 756-2584
1-800-422-9393
FAX (541) 756-5783
NORTH BEND, OREGON • 97459

JON C. KINTNER, M.D.
DENNIS E. OTTEMILLER, M.D.
DEBRA A. GRAHAM, M.D.
JANE GILBERT, M.D.
EYE M.D.s

Practice Limited to the Eye
Cataract Surgery & Intraocular Lenses
Board Certified, Members of American
Academy of Ophthalmology

How To Insert Eyedrops

Infections, inflammation, glaucoma, and many other eye disorders are treated with eye drops. Surprisingly, even the small amount of medication in an eye drop can create significant side effects in other parts of the body. It is important to remember that all medicines have potential side effects. There are ways to decrease the absorption of the eye drop into the system, and to increase the time the eye drop is on the eye, making the medicine more safe and effective.

Inserting eye drops may seem difficult at first but becomes easier with practice. To put in an eye drop, tilt the head back. Then create a pocket in front of the eye by pulling the lower lid down with an index finger or gently pinch the lower lid outward with the thumb and index finger. Let the drop fall into the pocket without touching your eye or eyelid (to prevent contamination of the bottle).

Immediately after instilling the drop, gently press back on the bridge of your nose for two to three minutes with your thumb and forefinger. This prevents most of the drop from traveling down the tear duct to the rest of the body.

Keep your eyes closed for three to five minutes after instilling the drop. Because the volume of a single drop exceeds the capacity of the surface of the eye, it serves no purpose to use two drops at the same time.